

الاسم :  
الرقم :مسابقة في الفيزياء  
المدة : ساعتان

*This exam is formed of three obligatory exercises in three pages*

*The use of non-programmable calculators is allowed*

**First Exercise ( 7 points) Suspension system in a car**

Certain tracks present periodic variations of its level .A car moves in a uniform motion on such a track that has regularly spaced bumps. The distance between two consecutive bumps is  $d$  and the speed of the car is  $V$ . In order to study the effect of the bumps on the car , we consider the car and the suspension system as a mechanical oscillator (elastic pendulum) whose oscillation takes a time  $T$ .

**A- Study of T**

**1. Theoretical study**

Consider a horizontal elastic pendulum formed of a solid of mass  $m$  attached to a spring of constant  $k$  and of negligible mass; the other end of the spring is fixed to a support. The forces of friction are supposed to be negligible and the solid of center of mass  $G$  can move on a horizontal axis  $Ox$ .

When the solid is at rest ,  $G$  coincides with the point  $O$  taken as origin of abscissa.

The solid is pulled from its equilibrium position by a distance  $x_m$ , and then released without initial velocity at the instant  $t_0 = 0$ . The horizontal plane passing through  $G$  is taken as a gravitational potential energy reference

At any instant  $t$  , the abscissa of  $G$  is  $x$  and the algebraic measure of its velocity is  $v$ .

- Starting from the expression of the mechanical energy of the system {pendulum -Earth}, determine the second order differential equation that characterizes the motion of the solid.
- Deduce the expression of its proper period  $T_0$  .

**2. Experimental study**

In order to show the effects of the mass  $m$  of the solid and the constant  $k$  of the spring on the duration of one oscillation of a horizontal elastic pendulum, we use four springs of different stiffnesses and four solids of different masses.

In each experiment , we measure the time  $\Delta t$  for 10 oscillations using a stopwatch .

**a) Effect of the mass  $m$  of the solid**

In a first experiment , the four solids are connected separately from the free end of the spring whose stiffness is  $k = 10$  N/m. The values of  $\Delta t$  are shown in the following table:

$m$ (g)	50	100	150	200
$\Delta t$ (s)	4.5	6.3	7.7	8.9

Determine, using the table, the ratio  $T^2 / m$ . Conclude.

**b) Effect of the stiffness  $k$  of the spring.**

In a second experiment , the solid of mass  $m = 100$  g is connected successively from the free end of each of the four springs. The new values of  $\Delta t$  are shown in the following table :

$k$ (N/m)	10	20	30	40
$\Delta t$ (s)	6.3	4.5	3.7	3.2

Determine, using the table , the values of the product  $T^2 \times k$ . Conclude.

### c) Expression of T

Deduce that T may be written in the form  $T = A\sqrt{\frac{m}{k}}$  where A is a constant.

### B) Oscillations of the car

- 1) The car, with the driver alone, forms a mechanical oscillator whose proper period is around 1s. It moves with a speed  $V = 36 \text{ km/h}$  on a path having equally spaced bumps. The distance between two consecutive bumps is  $d = 10 \text{ m}$ . The car enters then in resonance.
  - a) Specify the exciter and the resonator.
  - b) Explain why does the car enter resonance.
  - c) How can the driver avoid this resonance?
- 2) The driver, with four passengers, drives his car on the same path with the same speed of  $36 \text{ km/h}$ . Would the car enter in resonance? Justify your answer.

### Second Exercise (6 points) Energy levels of the hydrogen atom

The energies of the different energy levels of the hydrogen atom are given by the relation:

$$E_n = -\frac{13.6}{n^2} \text{ (in eV)} \quad \text{where } n \text{ is a positive whole number.}$$

Given :

$$\begin{aligned} \text{Planck's constant : } h &= 6.63 \times 10^{-34} \text{ J.s} & ; & & 1 \text{ eV} &= 1.60 \times 10^{-19} \text{ J} ; \\ \text{Speed of light in vacuum : } c &= 3 \times 10^8 \text{ m/s} & ; & & 1 \text{ nm} &= 10^{-9} \text{ m.} \end{aligned}$$

#### A- Energy of the hydrogen atom

- 1) The energies of the atom are quantized. Justify this using the expression of  $E_n$ .
- 2) Determine the energy of the hydrogen atom when it is:
  - a) in the fundamental state.
  - b) in the second excited state.
- 3) Give the name of the state for which the energy of the atom is zero.

#### B - Spectrum of the hydrogen atom

##### 1 - Emission spectrum

The Balmer's series of the hydrogen atom is the set of the radiations corresponding to the downward transitions to the level of  $n = 2$ .

The values of the wavelengths in vacuum of the visible radiations of this series are :

$$411 \text{ nm} ; 435 \text{ nm} ; 487 \text{ nm} ; 658 \text{ nm.}$$

- a) Specify, with justification, the wavelength  $\lambda_1$  of the visible radiation carrying the greatest energy.
- b) Determine the initial level of the transition giving the radiation of wavelength  $\lambda_1$ .
- c) Deduce the three initial levels corresponding to the emission of the other visible radiations.

##### 2 - Absorption spectrum

A beam of Sunlight crosses a gas formed mainly of hydrogen. The study of the absorption spectrum reveals the presence of dark spectral lines.

Give, with justification, the number of these lines and their corresponding wavelengths.

#### C - Interaction photon - hydrogen atom


- 1) We send on the hydrogen atom, being in the fundamental state, separately, two photons of respective energies  $3.4 \text{ eV}$  and  $10.2 \text{ eV}$ .

Specify, with justification, the photon that is absorbed.
- 2) A hydrogen atom found in its fundamental state absorbs a photon of energy  $14.6 \text{ eV}$ . The electron is thus ejected.
  - a. Justify the ejection of the electron.
  - b. Calculate, in eV, the kinetic energy of the ejected electron.

**Third Exercise ( 7 points) Saving life capacitor**

A heart suffering from disordered muscular contractions is treated by applying electric shocks using a convenient apparatus.

In order to study the functioning of this apparatus , we use a source of DC voltage of adjustable value  $E$  , a double switch , a resistor of resistance  $R$  and a capacitor ( initially neutral) of adjustable capacitance  $C$ . We connect the circuit represented in the adjacent figure.



**A. Theoretical study**

1. The switch is turned to position (1).

- a) Give the name of the physical phenomenon that takes place in the capacitor.
- b) Specify the values of the current in the circuit and the voltage  $u_{MN}$  after few seconds.

2. The switch is now turned to position (2) at an instant taken as  $t_0 = 0$ .

- a) Derive , at the instant  $t$  , the differential equation giving the variation of the voltage  $u_C = u_{MN}$  as a function of time.

- b) The expression  $u_C = A e^{-\frac{t}{\tau}}$ , where  $A$  and  $\tau$  are constants , is a solution of that equation.

Determine the expressions of  $A$  and  $\tau$  in terms of  $E$ ,  $R$  and  $C$ .

- c) Derive the expression giving the current  $i$  during the discharging as a function of time.


**B. Using the apparatus**

The energy needed to save the life of a patient during an electric shock is 360 J. This energy is supplied by discharging the capacitor through the patient's chest (ribcage) considered as a resistor of resistance  $50 \Omega$  during a time  $t_1$  that can be controlled by the switch.

The capacitance of the capacitor is adjusted on the value

$C = 1$  millifarad and is charged under the voltage

$E = 1810$  V.



1) Determine the energy stored in the capacitor at the end of the charging process.

2) The discharging starts at the instant  $t_0 = 0$  .At the instant  $t_1$  , the energy delivered to the patient amounts to 360J ,the switch is then opened .

- a) Calculate the energy that remains in the capacitor at the instant  $t_1$ .
- b) Using the results of the above theoretical study; determine:
  - i) the value of  $t_1$  .
  - ii) the current at the end of the electric shock.

## First Exercise

**A) 1 - a)** M.E =  $\frac{1}{2}kx^2 + \frac{1}{2}mv^2$  ;

No friction the M.E is conserved  $\Rightarrow \frac{dM.E}{dt} = 0 \Rightarrow kxv + mvx'' = 0$

$\Rightarrow x'' + \frac{k}{m}x = 0$

b)  $\omega_0^2 = \frac{k}{m} \Rightarrow \omega_0 = \sqrt{\frac{k}{m}}$  ;  $T_0 = \frac{2\pi}{\omega_0}$

$\Rightarrow T_0 = 2\pi\sqrt{\frac{m}{k}}$

**2 - a)**  $\frac{T^2}{m} = 4$  (S.I)  $\Rightarrow \frac{T^2}{m} = \text{constant}$

b)  $T^2 \times k = 4$  (S.I)  $\Rightarrow T^2 \times k = \text{constant}$

c) T is proportional  $\sqrt{m}$  to and T is inversely proportional to  $\sqrt{k}$

$\Rightarrow T = A\sqrt{\frac{m}{k}}$

**B) 1 - a)** Exciter is the bumps and the resonator is the car

b) The car is submitted to pulses

periodically of period :  $T' = \frac{d}{V} = 1\text{sec}$

$T_0 = 1 \text{ sec} ; T' = T_0 \Rightarrow \text{Resonance}$

c) Mass increases  $\Rightarrow T_0$  increases

$\Rightarrow T_0 \neq T'$

## **Second Exercise**

**A) 1** –  $E_1 = -13.6 \text{ eV}$  ;  $E_2 = -3.4 \text{ eV}$  ;  $E_3 = -1.51 \text{ eV}$  ;  $E_\infty = 0$

$\Rightarrow$  The values of energies are discontinuous .

**2** – a)  $E_{\text{fund.}}$  corresponding to  $n = 1 \Rightarrow E_{\text{fund.}} = -13.6 \text{ eV}$

b) Second excited state corresponding to  $n = 3$

$\Rightarrow E_3 = -1.51 \text{ eV}$ .

**3** – Ionize state

**B) 1** – a)  $E = \frac{hc}{\lambda}$  or  $E$  is inversely prop. to  $\lambda$

$\Rightarrow \lambda_1 = 411 \text{ nm}$

b)  $\frac{hc}{\lambda} = E_i - E_f \Rightarrow \frac{hc}{\lambda} = \left( \frac{-13.6}{n^2} + \frac{13.6}{4} \right) 1.6 \times 10^{-19} \text{ J}$  ;

For  $\lambda = \lambda_1$  ;  $n = 6$

c) The other three levels are :  $n = 5$  ;  $n = 4$  ;  $n = 3$  to  $n = 2$

**2** – The dark lines of the absorption spectrum corresponding to the bright lines of same wavelength of the emission spectrum .

We have 4 bright lines  $\Rightarrow$  we have 4 dark lines of wavelengths : 411 nm ; 487 nm ; 658 nm

**C) 1** –  $-13.6 + 3.4 = -10.2 = \frac{-13.6}{n^2} \Rightarrow n = 1.15$  ;

$n$  is not a whole number  $\Rightarrow$  not absorbed

$-13.6 + 10.2 = -3.4 = \frac{-13.6}{n^2} \Rightarrow n = 2$  (whole no)  $\Rightarrow$  absorbed

**2** - a) The energy of the photon is greater than the ionization energy

b)  $\text{K.E} = -13.6 + 14.6 = 1 \text{ eV}$

c)

### Third Exercise

**A) 1** – a) Charging of the capacitor

b)  $i = 0$  ;  $u_C = E$  .

**2** – a)  $u_C = Ri = -RC \frac{du_C}{dt}$

$$\Rightarrow u_C + RC \frac{du_C}{dt} = 0$$

b) At  $t = 0$  ;  $u_C = A = E$  ; Derive  $u_C$  and substitute  $\Rightarrow \tau = RC$

c)  $i = -C \frac{du_C}{dt} \Rightarrow i = \frac{E}{R} e^{-\frac{t}{\tau}}$

**B) 1** –  $E = \frac{1}{2} CU^2 \Rightarrow E = 1638 \text{ J}$

**2** – a)  $E_{\text{rem.}} = 1638 - 360 = 1278 \text{ J}$

b) i)  $E_{\text{rem.}} = \frac{1}{2} C u_C^2$

$$\Rightarrow u_C = 1599 \text{ V ;}$$

$$u_C = E e^{-\frac{t}{\tau}} \Rightarrow t = 6.2 \text{ ms}$$

ii)  $i = \frac{E}{R} e^{-\frac{t}{\tau}} \Rightarrow i = 32 \text{ A}$